

Quantifying dry matter production and partitioning in different parts of safflower (*Carthamus tinctorius* L.)

*B. Torabi¹, H. Saadatkhan², A. Soltani³ and B. Mahdavi⁴

¹Assistant Prof., Dept., of Agronomy, Gorgan University of Agricultural Sciences and Natural Resources,

²M.Sc. Student, Dept., of Agronomy, Vali-e- Asr University of Rafsanjan,

³Professor, Dept., of Agronomy, Gorgan University of Agricultural Sciences and Natural Resources,

⁴Assistant Prof., Dept., of Agronomy, Vali-e- Asr University of Rafsanjan

Received: 09/06/2015; Accepted: 05/10/2016

Abstracts

Introduction: Simulation crop models are a robust tool to improve crop management and to study yield limiting and reducing factors. Crop parameters related to phenology, leaf area and dry matter production and partitioning are needed to simulate crop growth and yield. Therefore, the aim of present study is to quantify the dry matter production and to estimate dry matter partitioning coefficients to different crop parts (stem, leaf and grain).

Materials and Methods: A factorial experiment was conducted based on completely randomized block design with four replicates and three sowing dates (4 April, 25 April and 16 May) and four cultivars (411, Sina, Local Isfahan and Sofeh) in Research Farm of Rafsanjan Vali-e-Asr University in 2011. Sampling was done at an interval of 5-10 days from two weeks after planting and continued up to end of the growing season. On each sampling, the leaf, stem and grain dry matter weight were measured. The truncated exponential model was fitted on total dry matter weight (w) data over the time (t). Coefficient of dry matter partitioning to different crop parts (leaf, stem and grain) was obtained by fitting the linear regression model on dry matter weight of each part versus total dry matter weight.

Results: Results showed that the model well described the trend of dry matter production versus days after planting. According to the model, the maximum dry matter accumulation in all three planting dates was obtained 972-1179, 576-611 and 191-277 gm^{-2} , respectively. Coefficient of dry matter partitioning to leaves ranged from 0.45 to 0.51 in the first planting date, 0.51 to 0.60 in the second planting date and 0.44 to 0.61 in the third planting date. There was no different between the coefficients of cultivars. Coefficient of dry matter partitioning to stem in all three planting dates were 0.35-0.49, 0.24-0.44 and 0.14-0.23, respectively. There was no significant difference among the cultivars. Partitioning coefficient to grain was between 0.16 and 0.44 for the first planting date that was lower than the second planting date. In the third planting date, the coefficient was between 0.35 and 0.76 that there was a significant difference among the cultivars.

Conclusion: It was concluded that the trend of the dry matter production and partitioning is affected by environmental condition (temperature and solar radiation) and cultivar. The results showed that with delay in planting date crop growth rate and dry matter production were decreased. In such condition, coefficient of dry matter partitioning to stem was decreased and coefficient of dry matter partitioning to leaves and grain was increased.

Keywords: Dry matter, Planting date, Quantification, Safflower

*Corresponding author: Ben-Torabi@yahoo.com

Interaction between sulfur and zinc on cotton (*Gossypium sp. L.*) yield and some soil parameters

Kh. Sori Abdoulahzadeh¹, *Sh. Shahsavani², H. Abasdokht³, A. Abaspoor⁴ and
M. Rahimi⁵

¹M.Sc. Student of Soil Science, Shahrood University, ²Assistant Prof., Dept., of Soil Science,
Shahrood University, ³Associate Prof., Dept., of Agronomy and Plant Breeding, Shahrood University,

⁴Associate Prof., Dept., of Soil Science, Shahrood University, ⁵Lecturer, Dept., of Agronomy and
Plant Breeding, Shahrood University

Received: 17/04/2016; Accepted: 02/05/2017

Abstract

Introduction: Cotton cultivation from income and employment point of view in the world and developing countries is very important. Cotton cultivation in the world in the beginning of the century was 35 million hectare and its yield was 53 million tone. The highest production was belonging to countries like China, America and India. Chemical fertilizers especially in past, contained sulphur, especially simple super phosphate, ammonium sulphate, potassium sulphate and mixed fertilizers which contain excess sulphur and until using these fertilizers sulphur deficiency will not appear in the farm, but from economical point of view farmers prefer to use more refine fertilizers with low volume. In warm and semi-warm area of the world sulphur used as a plant nutrition mineral as its deficiency continually reported (Falahatgar et al., 2013). Sulphur has important role in plant producing different kinds of amino acids like methionine and cysteine (Asgha Malik et al., 2004). Sulphur in soil is under influence of *Thiobacillus* bacteria which convert it to sulphuric acid in the soil and this sulphuric acid can affect the soil reaction (Dawood et al., 1985).

Method and materials: This experiment was conducted in the form of factorial on the basis of complete randomized block design with 12 treatments and three replications with 36 plots. The factors included the using, zinc sulphate in two levels, granular sulphate in three levels, control, 500 and 1000 kg ha⁻¹ and using *Thiobacillus* in two levels, control and application. After harvesting soil sampling and plant leaf collection were done from each plot. Each plot had 5 cultivated row with three meter length and distance between two rows were 75cm, plants distance were 20 cm. Granular sulphur and *Thiobacillus* were applied to soil before sowing and all basal fertilizers were applied at the time of sowing only urea were applied in three split, one third at the time of sowing, one third at tillering and last one at flowering. At different time of growing like appearance of 4 leaves or 8 leaves thinning and weeding were done. During growing season some agronomic properties like cotton yield, cotton seed yield, mean boll weight, plant boll number, reproductive branch number, vegetative branch number, plant height and cotton kale percent. Also after harvesting soil sampling were done from each plot for soil properties analysis like Ec, pH, total nitrogen, available phosphorus and potassium, iron, zinc etc. all data were analyzed for analysis of variance using MSTATC and mean comparison were done for significant difference (LSD) at 5% level.

Results and Discussion: Results of analysis of variance data showed that, sulphur effect and its interaction effect with *Thiobacillus* and zinc were significant at 5% level. Results of mean comparison of sulphur effect and its interaction with *Thiobacillus* and zinc on cotton yield

*Corresponding author: shahsavani2001@yahoo.com

showed that treatment with 1000 kg ha⁻¹ sulphur without zinc and *Thiobacillus* were 5760 kg ha⁻¹ and with *Thiobacillus* and zinc cotton yield were 5616 kg ha⁻¹, and with zinc and without *Thiobacillus* were 5853 kg ha⁻¹. Results of analysis of variance for cotton seed yield showed that sulphur effect and zinc and its interactions with *Thiobacillus* and zinc were significant at 1% level and interaction between sulphur and zinc on cotton seed yield were significant at 5% level. Results of analysis of variance on fiber yield showed that interaction effects of sulphur and *Thiobacillus* on fiber were significant at 1% level, but interaction effect of zinc and *Thiobacillus* were significant at 5% level. Results of analysis of variance showed that effect of sulphur on kale percent were significant at 1% level and its interaction with *Thiobacillus* on kale percent were significant at 5% level. Results of analysis of variance showed that sulphur effect and its interaction with *Thiobacillus* and interaction effect of zinc and *Thiobacillus* on boll weight were significant at 1% level, but effect of zinc and *Thiobacillus* on boll weight were significant at 5% level. Results of analysis of variance showed that sulphur effect and its interaction with zinc and *Thiobacillus* on reproductive branches were significant at 1% level. Results of analysis showed that effect of zinc and its interaction with sulphur on reproductive branches were significant at 1% level. Sulphur and *Thiobacillus* effect on vegetative branches were significant at 5% level. Results of analysis of variance showed that sulphur effect and zinc and the interaction effect of sulphur and *Thiobacillus* on number of bolls in plant were significant at 1% level also main effect of sulphur, zinc and *Thiobacillus* and their interaction with zinc on oil percent were not significant.

Conclusion: Results showed that interaction effect of sulphur, zinc and *Thiobacillus* on some agronomical properties and soil were significant. According to mean comparison results, treatment of 1000 kg ha⁻¹ sulphur, without *Thiobacillus* and zinc caused significant effect on cotton yield, cotton seed yield, soil Ec, soil available sulphate and zinc, also application of 1000 kg ha⁻¹ sulphur with *Thiobacillus* increased cotton yield and cotton seed yield compare to control.

Keywords: Cotton, Sulphur, Zinc

Phosphorus and dry matter accumulation and partitioning coefficients as affected by fertilizer phosphorus rate and inoculation by *Streptomyces* bacteria in wheat and barley

T. Khosravian¹, *E. Zeinali², A. Siahmarguee², R. Ghorbani Nasrabadi² and S.M. Alimaghani¹

¹Ph.D. Student, Gorgan University of Agricultural Science and Natural Resources,

²Faculty of Member, Gorgan University of Agricultural Science and Natural Resources

Received: 15/08/2016; Accepted: 09/11/2016

Abstract

Background and objectives: Information on the concentration and accumulation of phosphorus (P) in the plant dry matter (DM) and the coefficients of its allocation to different plant parts can help improve the P nutrition management of crops and increase the efficiency of P fertilizers. In addition, this information is required in crop growth simulation models.

Materials and methods: In order to investigate the effect of inoculation with *Streptomyces* sp. bacteria and fertilizer P rates (0, 20, 40, 60 and 80 kg P ha) on the P and DM accumulation and partitioning in wheat (cultivar Morvarid) and barley (cultivar Sahra) a pot experiment was carried out in new campus of Gorgan University of Agricultural Sciences during 2015 growing season. The experimental design was completely randomized design as a factorial experiment with four replications. Soil available P was recorded 5.8 mg kg⁻¹.

Results: Results showed that the effect of fertilizer P rate on DM accumulation, P concentration and accumulation in all plant parts was significant ($P = 0.01$), statistically. Inoculation also influenced on the DM accumulation in all plant parts except seeds, and P accumulation in all of them except the stem, but its effect on the P concentration in different parts of plant with the exception of leaf and root was not significant. Hence, it can be deduced that the rate of fertilizer phosphorus through both DM accumulation and P concentration, and inoculation only via DM accumulation affected on the P accumulation. Also, the findings of this study showed a very high stability of allocation coefficients of DM and P to different parts of plant compared to DM and P accumulation and P concentration. In unfertilized control, the accumulated phosphorus rate in whole plant was 3.99 and 4.57 and in grain was 1.79 and 1.94 mg P plant⁻¹ for wheat and barley, respectively. Based on the segmented model fitted data, the accumulation of phosphorus in the whole plant increased linearly up to 51 and 59, and in their grain up to 47 and 54 Kg P ha⁻¹ for wheat and barley, respectively. But after that, more fertilizer P application did not significantly change the quantity of P accumulated per plant or grain.

In above mentioned fertilizer phosphorus rates, the amount of P accumulated in whole plant was 13.67 and 13.38 and in grain was 7.95 and 6.29 mg P plant⁻¹ for wheat and barley respectively. In both investigated crops, at all levels of fertilizer P, the order of organs based on the P accumulation was as grain > stem > leaf > root.

Conclusion: The findings indicate the increased dry matter accumulation, concentration and accumulation of phosphorus in all plant organs by increasing the amount of phosphorus fertilizer, and relative stability of allocation coefficient of dry matter and phosphorus to organs at different levels of phosphorus fertilizer. Also, inoculation increased vegetative growth and thereby increased the accumulation of phosphorus in plant. The results of this research reveals that two strategic crops of wheat and barley uptake 3.6 to 5.9 kg P per ton of grain and 1.7 to 2.9 kg P per ton of straw depending on the amount of available P in the soil.

Keywords: Grain, Leaf, Phosphorus concentration, Root, stem

*Corresponding author: e.zeinali@yahoo.com

Effects of different levels of vermicompost on morphophysiological and essential oil characteristics of Peppermint (*Mentha piperita* L.) under water deficit regimes

*H. Gorgini Shabankareh¹ and S. Khorasaninejad²

¹Ph.D. Student, Dept., of Horticulture, Gorgan University of Agricultural Sciences and Natural Resources, Iran, ²Assistant Prof., Dept., of Horticulture, Gorgan University of Agricultural Sciences and Natural Resources, Iran

Received: 22/08/2016; Accepted: 02/05/2017

Abstract

Background and objectives: Drought is one of the most important factors limiting plant growth around the world. So it can be used minimum water with using natural fertilizer such as vermicompost and can be caused to reduce water evaporation from soil, protect and storage water in the soil. The interest in plant products has increased considerably all over the world because many herbal medicines are free from side effects since most medicinal plants are consumed raw, proper management of crop production is needed to achieve high quality plants. So in this regard, experiment was done to study the effect of different levels of vermicompost on morphophysiological and essential oil characteristics of Peppermint (*Mentha piperita* L.) under water deficit regimes.

Materials and methods: A pot experiment was conducted as factorial based on completely randomized design with three replications at Gorgan University of Agricultural Sciences and Natural Resources greenhouse during 2016. Treatments were included Irrigation regimes in three levels (100 percent FC, control, 75percent FC and 50percent FC) and vermicompost in three levels (0, 20 and 40percent volumetric). Studied traits were included plant height, internodes distance, fresh and dry weight, number of tillers, leaf water ratio, proline, soluble sugar and essential oil percent. After planting, harvesting and growth characteristics, essential oil, soluble sugar and proline were evaluated with Clevenger, the method of Omokolo and the method of Bates, respectively.

Results: The results showed that internodes distance, fresh and dry weight of shoot, number of tiller and leaf water ratio, were decreased by increasing drought stress. Also, with increasing drought stress, leaf water ratio decreased (2.13 percent) and prolin, soluble sugar and essential oil percent were increased 0.72, 10, 14 and 0.13 percent, respectively. Vermicompost had significant effects on all of morphological and physiological characteristic and this effect was the highest in 40 percent volumetric. Interaction effects of Irrigation regime and vermicompost had significant effects on all of morphological and physiological characteristic except of fresh weight of shoot and number of tillers. Essential oil (0.33 percent) and leaf water ratio (23.71 percent) increased in 40 percent volumetric in 50 percent FC.

Conclusion: Due to the results, it be expressed to reduce chemical fertilities use and achieve to sustainable agriculture, can be adjusted with vermicompost application a lot of part from drought stress effects on Peppermint. Generally, the results showed that by vermicompost application (40 percent volumetric) reduced drought damage on growth characteristics and the most essential oil.

Keywords: Essential oil, Lamiaceae, Proline, Vermicompost, Water deficit stress

*Corresponding author: h.shabankareh92@gmail.com

Sprouting, plant establishment, and yield improvement of potato (*Solanum tuberosum* L.) minituber cultivars by foliar application of benzylaminopurine and abscisic acid

M.J. Ahmadi Lahijani¹, *M. Kafi², A. Nezami², J. Nabati³ and J. Erwin⁴

¹Ph.D. Dept., of Agronomy and Plant Breeding, Ferdowsi University of Mashhad, Iran,

²Professor, Dept., of Agronomy and Plant Breeding, Ferdowsi University of Mashhad, Iran,

³Assistant Prof., Research Center of Plant Sciences, Ferdowsi University of Mashhad, Iran,

⁴Professor, Dept., of Horticultural Science, University of Minnesota, USA

Received: 30/10/2016; Accepted: 04/02/2017

Abstract

Backgrounds and objectives: Potato is the fourth important crop worldwide. Since potato is mainly propagated vegetatively via tubers, increasing production of high quality and virus free seed potato is of great importance. Phyto-hormones regulate a wide range of processes and tuberization in potato. Due to small size and weakness of minitubers and some difficulties in plantlets establishment in the field, more studies are needed to improve plant establishment and productivity of minitubers under such conditions. Therefore, this study was carried out in order to evaluate the possibility of improving sprouting and plantlet establishment of potato minituber cultivars (Agria and Fontane) using plant growth regulators (6-benzylaminopurine- BAP, and abscisic acid- ABA).

Materials and Methods: A two-step experiment was carried out at the Ferdowsi University of Mashhad, during 2014-15. In the first experiment, foliar treatment of mother plants at stolonization or tuberization stages using plant growth regulators (BAP, ABA, and BAP+ABA, 50 μ M) under controlled conditions was performed. In the second experiment, progeny minitubers were compared in terms of sprouting, plantlets establishment, and yield under field conditions. Factorial arrangement based on completely randomized design and complete block design were used for the first and second experiments, respectively. Number of sprouts per minituber, length of sprouts, day to emergence, emergence percentage, number of stems per plant, morphological traits, yield, and yield component were evaluated.

Results: Day to emergence tended to decrease with an increase in sprout length. In both cultivars, BAP application at tuberization stage increased sprout length compared with the untreated control plants. The number of established plantlets and percentage of emergence were enhanced by treating plants using BAP or ABA compared with control plants, but the effects were different between cultivars. The greatest emergence percentage observed in ABA-treated plants at stolonization stage, which was 39% more than the control. In addition, Fontane showed 15% higher emergence rate than Agria. BAP+ABA application at tuberization stage advanced the number of sprouts in Agria, whereas, the most number of sprouts in Fontane recorded when BAP applied at tuberization stage. Good linear relationship was observed between sprout length and day to emergence ($R^2=0.88$) and between the number of sprouts on minitubers and the number of tubers in the next generation ($R^2=0.53$). Regardless to the cultivar, application of BAP+ABA at tuberization stage increased the number of tubers per plant.

Conclusion: The results indicating that application of BAP resulted in increasing the number of sprouts and shortened the days to emergence, whereas, ABA improved emergence parentage and plantlet establishment of the both cultivars. It seems that both BAP and ABA could be effectively used for improving sink strength for better supporting the sprouts growth under field conditions. Based on our results, application of PGRs increased the number of sprouts per tuber which result in producing higher number of stems, hence, enhance the number of tubers per plant, however, the effects are different between cultivars.

Keywords: Cytokinin, Days to emergence, Sprout length, Plant Growth Regulators

*Corresponding author: m.kafi@um.ac.ir

Early sowing date as a strategy for improvement of maize yield and maize physiological and phenological characteristics in climate change conditions at Kermanshah Province

*S. Rahimi-Moghaddam¹ and Kh. Azizi²

¹Ph.D. Graduate, Dept., of Agroecology, Shahid Beheshti University,

²Associate Prof., Dept., of Agronomy and Plants Breeding, Lorestan University

Received: 31/10/2016; Accepted: 12/06/2017

Abstract

Background and objectives: Climate change can directly affect on worldwide food security because climate change processes (such as increasing of CO₂ concentration and temperature and variability of precipitation) have directly affect on crops. Corn such as C₄ plants that are sensitive to climate changes. Therefore to reduce the sensitivity of the maize in the face of climate change, we need to apply adaptation strategies. One of the effective strategies is the changes in sowing dates. Many of studies have shown that changes in sowing dates (particularly the use of early planting dates) can reduce the negative effects of climate change.

Materials and methods: This research was conducted in three locations of Kermanshah Province. Accordingly, the future climate in the study areas was generated using long-term (1980-2009) climate data of the baseline (included minimum and maximum temperatures, rainfall and global radiation) and AgMIP technique under two climate scenarios (RCP4.5 and RCP8.5) for the future period of 2040 -2069. Long-term simulation experiments consisted of five sowing dates (5st April, 20st April, 5st May, 20st May, 5st June), three locations (Kermanshah, Kangavar and Eslamabad), two future climate scenarios (RCP4.5 and RCP8.5) in 30 years. In total, around 1350 simulation experiments were carried out. In this study, APSIM crop model was used for simulation of maize growth and yield. In this study, all of the simulations were conducted in potential conditions and water- and nitrogen-limited production situations were not considered in the current study. APSIM model has previously been calibrated and evaluated for SC704 cultivar (this cultivar is the most common cultivated cultivar in the Kermanshah Province). In current research, all the output data was analyzed, graphed and mapped using the R software package (24) and OriginPro9.1 (31).

Results: Average grain yield of Kermanshah Province was 11354 kg ha⁻¹ in the baseline. Results showed that in 2050, on average grain yield was reduced 60.82 and 80.73 % (under RCP4.5 and RCP8.5, respectively) compared to baseline. In different locations, the highest and lowest grain yield in the baseline were recorded in the Kangavar with 13426 kg ha⁻¹ and Kermanshah with 7952.4 kg ha⁻¹. When averaged across locations, the highest grain yield in the climate change conditions was obtained in an early sowing date (5st April) with 7071.2 and 4743 kg ha⁻¹ (under RCP4.5 and RCP8.5, respectively). In future on average (two scenarios) duration of growth period, vegetative and reproductive growth periods (4.7, 4 and 1.7 %, respectively) were decreased compared to the baseline. Also, in future, the number of grains and grain weight were reduced in Kermanshah Province so that under RCP4.5 were 56.5 and 31.8 %, respectively and under RCP8.5 were 78.5 and 59.3 %, respectively. However, these reductions in duration of growth period, vegetative and reproductive growth periods, the number of grains and grain weight were less in 5st April early sowing date than other sowing dates (especially late sowing dates).

Conclusion: Generally, results of the current study indicated that climate change had negative effects on maize yield and maize physiological and phenological characteristics in Kermanshah Province. However, early sowing dates might reduce these negative effects. So that in most cases, 5st April can reduce negative effects on maize yield and maize physiological and phenological characteristics in the future period.

Keywords: APSIM-Maize, Climate change, Number of grains, Maximum leaf area index

*Corresponding author: sajadr.moghaddam@yahoo.com

Responses of some agronomical characteristics of faba bean (*Vicia faba* L.) to biofertilizers under water deficit stress

*E. Rezaei Chiyaneh¹, A. Rahimi¹, F. Sheikh² and M. Mohayeji³

¹Assistant Prof., Dept., of Agronomy, Urmia University, Iran, ²Assistant Prof., Golestan Agricultural and Natural Resources Research and Education Center, ³Agricultural Research, Education and Extension Organization, Gorgan, Iran, ⁴Assistant Prof., Dept., of Agronomy and Plant Breeding, Shahid Bahonar University of Kerman, Iran

Accepted: 23/11/2016; Received: 12/04/2017

Abstract

Background and objectives: Iran's climate is mostly arid and semi-arid, where water availability is a major problem in crop production. In regions where water scarcity is the principle limiting factor for crop cultivation, farmers are interested in growing crops that are able to adapt to drought conditions. Bio-fertilizer has been identified as an alternative to chemical fertilizer in sustainable farming. Organisms that are commonly used as biofertilizers component are nitrogen fixers (N-fixer), potassium solubilizer (K-solubilizer) and phosphorus solubilizer (P-solubilizer), or with the combination of molds or fungi increased soil fertility, growth and yield plants. Therefore, the aim of this study was to investigate the use of biofertilizer in drought stress on several morphological and yield characteristics of faba bean under field condition.

Materials and methods: In order to study the effects of different irrigation levels and biofertilizers on yield and yield components of faba bean, a field experiment was conducted as split plot based on a Randomized Complete Block Design with three replications at a farm located in West Azerbaijan province – city of Naqadeh, during growing season of 2013-2014. The main plots were allocated to irrigation levels (Irrigation after 50 mm (control), 100 mm (moderate water stresses) and 150 mm (severe water stresses) evaporation from class A pan) and sub-plots were allotted to biofertilizers (*Azotobacter* and Barvar Phosphate-2, *Azotobacter* + Barvar Phosphate-2 and control). The experimental units in each block included of 8 rows with 40 cm inter-row and 7 cm intra-row spacing of 4 meters long. Different traits such as plant height, number of pods per plant, number of seeds per pod, 1000-seed weight, biological yield, seed yield and percentage of protein were recorded. Seed protein content was determined by the micro-Kjeldahl method. Seed protein content was calculated by multiplying the total N content by 6.25. The data were subjected to analysis of variance (ANOVA) using SAS statistical package. Means of each trait were compared on the bases of LSD test at $p \leq 0/05$.

Results: Results showed that factors had significant effects on plant height, number of pods per plant, number of seeds per pod, 1000-seed weight, biological yield and seed yield of faba bean but the interaction of irrigation×biofertilizer on any of the traits under study were not significant. In comparison of severe stress (lowest yield and yield components) and non-stress conditions, plant height, number of pods per plant, number of seeds per pod, 1000-seed weight, biological yield and seed yield were decreased 21.90, 39.83, 28.83, 26.15, 33 and 38%, respectively. Increasing of distance irrigation increased grain protein content. Also, the use of bio-fertilizers increased the protein content of faba bean. Generally, combined usage of biofertilizers showed the greater increasing in studied traits than individual consumption.

Conclusion: Inoculation with biofertilizers could not reduce the adverse effects of drought stress and could not effect on increasing of yield significantly. Therefore, to achieve optimal performance in arid and semi-arid areas, it's better to irrigation faba bean during different periods of growth and developmen. It is suggested that this research is better to be conducted on another year to get the best functioning for this area.

Keywords: *Azotobacter*, Barvar Phosphate-2, Pod, Yield components, Yield

*Corresponding author: e.rezaeichiyaneh@urmia.ac.ir

Response of different genotypes of safflower (*Carthamus tinctorius* L.) to the foliar spraying of nano-iron oxide at the low irrigation conditions

S. Parsamehr¹, *R. Aminian² and F. Habibzadeh³

¹M.Sc. Student, Dept., of Production and Plant Breeding, International University of Imam Khomeini, Qazvin, ²Assistant Prof., Dept., of Production and Plant Breeding, International University of Imam Khomeini, Qazvin

Accepted: 06/01/2017; Received: 01/07/2017

Abstract

Background and objectives: Among oilseeds that have a good compatibility with the climatic conditions of Iran, safflower has a special place due to the drought and salinity resistance and also because of having the different spring and autumn varieties. Environmental stresses are the main factors reducing growth and crop yield. Among the environmental stresses, water stress is the main risks for the successful production of crops in Iran and the world. One of the most important applications of nanotechnology in various agricultural fields, especially regarding to the water and soil, is the application of nano-fertilizers to nourish plants. Iron is one the micro elements in plant nutrition. In the condition of iron deficiency, the amount of photosynthetic pigments such as chlorophyll content decreases.

Materials and methods: To evaluate the effect of foliar spraying of nano-iron oxide on yield and some important agronomic traits of safflower genotypes at the low water stress conditions, an experiment was carried out in split factorial based on randomized complete block design with three replications in 2016 at the research farm of Imam Khomeini International University. The main factors include irrigation on two levels (full irrigation and water stress in the flowering stage) and nano-iron oxide spraying on two levels (application and non-application of nano-iron oxide) and sub-factor included 10 genotypes of safflower (Golmehr, Padideh, Goldasht, Sofeh, Mexican88, Sina, Mexican11, Isfahan mahali, Faraman and Kuseh).

Results: The effect of irrigation on plant height, seed number in boll, boll number in plant and seed yield were significant. The effect of fertilizer was significant only on the subsidiary branches number but genotype effect was significant on all traits. The interactions between irrigation and genotype, fertilizers and genotype and among irrigation, fertilizer and genotype were significant on all traits. The interaction between irrigation and fertilizer in most of the traits except 1000 grain weight and harvest index was significant. Mexican88 genotype had the maximum grain yield (499.5 gr. m²) in the conditions of non-stress and foliar application of nano-iron oxide. Correlation analysis revealed that harvest index in both normal and stress conditions showed the highest correlation with grain yield. Based on cluster analysis, Sina, Kuseh, Goldasht and Mexican 88 genotypes in non-stress condition and Faraman, Kuseh and Mexican 88 genotypes under stress condition had the highest yield and yield components.

Conclusion: The study showed that, regarding to the most traits, Mexican 88 and Kuseh genotypes were suitable for planting in both stress and non-stress conditions.

Keywords: Water deficit stress, Nano-iron oxide, Safflower, Yield

*Corresponding author: roghayehaminian@yahoo.com

Determination of optimum concentration of three antioxidant enzymes for increased drought tolerance in Mung Bean (*Vigna radiata* L.) using genetic algorithm

Sh. Skandranejad¹, *M. Gholipoor² and H. Makarian²

¹Ph.D. Student, Dept., of Agronomy, Shahrood University of Technology, Shahrood, Iran,

²Associate Prof., Dept., of Agronomy, Shahrood University of Technology, Shahrood, Iran

Accepted: 07/01/2017; Received: 12/06/2017

Abstract

Background and objectives: The free radicals amounts are increased due to drought stress and subsequently they harm plant. The drought stress-resulted free radicals are scavenged by changing in activity of antioxidants like catalase, superoxide dismutase (SOD) and guaiacol peroxidase (GP) (regressors or independent variables; Xs). This reaction is known as resistance to (protection against) oxidative stress. This protection is reflected in traits like growth and grain production (biomass and harvest index, respectively; dependent or response variables; Ys). By maximizing Ys in relation to Xs, the combination of Xs with possible highest Ys can be obtained. This optimization comes from the relations of Ys with Xs under drought stress conditions. Due to having more than one Ys here, the relation tends to be multivariate and highly complex, especially when there is no strong positive relation (correlation greater than +0.95) between Ys variables. In such situations, the genetic algorithm can overcome the complexity. The output of mentioned algorithm can be used by plant breeders. In another words, breeders can increase drought tolerance by genetic manipulation of plant for optimum activity of Xs. This experiment was aimed to do the mentioned optimization for Mung Bean.

Materials and methods: The experiment was carried out in pots located in an open filed to increase the accuracy and possibility of generalizing the results to field results. Pots had 5 kg capacity in which 5 seeds of line VC1973a were planted. For thinning, 3 seedlings were removed, and left 2 ones. The filed capacity was determined using weight method. Treatment levels were 4 levels of low irrigation including irrigation at 80% (control), 65%, 50%, and 35% of field capacity. At maturity stage, the harvest index and biomass (biological yield) (Ys) were measured. The concentration of antioxidant enzymes catalase, SOD, and GP (Xs) were determined at flowering stage. For maximizing the function, first partial desirability function was determined. Then general desirability function was calculated. The value of Xs for which the highest amount of Ys is attainable was obtained on the basis of genetic algorithm and using MATLAB software.

Results: The results indicated that a function with 7 components including the main and interactive effects of Xs could predict the relation of Ys with Xs well (Adjusted $R^2 \geq 0.97$). The standardized regression coefficient was positive for catalase which reveals that the drought resistance (harvest index and biomass) enhances with increasing the activity of this enzyme. Like catalase, the effect of GP was additive on drought tolerance, but considering its standardized regression coefficient, this enzyme had a lower effect than catalase. Due to negative standardized regression coefficient for SOD, it could be concluded that the drought tolerance doesn't enhance with increasing activity of SOD.

Conclusion: The optimized concentration of catalase, SOD, and GP was $0.956 \mu\text{mol H}_2\text{O}_2 \text{ g}^{-1} \text{ FW}$, $24.23 \text{ AU g}^{-1} \text{ FW}$, and $21.23 \text{ AU g}^{-1} \text{ FW}$, respectively for possible maximum drought tolerance. It should be mentioned that these optimized concentrations were all among observed concentrations. Moreover, reports indicate that there is genetic diversity in activity of antioxidant enzymes (prerequisite to carry out breeding programs for attaining the optimized activity) for Mung Bean.

Keywords: Optimization, Catalase, Superoxide dismutase, Guaiacol peroxidase

*Corresponding author: manouchehr.gholipoor@gmail.com

The effect of folic acid on yield and some qualities parameters of wheat

A. Javadi¹ and *E. Esfandiari²

¹Ph.D. Graduated of Crop Physiology, Dept., of Agronomy and Plant Breeding, University of Maragheh,

²Professor, Dept., of Agronomy and Plant Breeding, University of Maragheh

Received: 09/02/2017; Accepted: 03/12/2017

Abstract

Background and purpose: Nowadays malnutrition due to low food diversity is a serious problem in developing countries like Iran. Due to important role of wheat in nutrition of these countries, researchers try to produce wheat grains with higher proteins, amino acids, iron and zinc content, providing essential body nutrients and as a result reduce malnutrition. Considering the role of folic acid in transfer of single carbon fragments and its effect on improvement of metabolic processes, in this research effect of external application of folic acid on the quality and yield of wheat grains were investigated.

Material and methods: Gaskogen wheat cultivar was planted in a farm experiment conducted at research farm of Maragheh University in early spring of 2013 using randomized complete block design with three treatments with density of 500 seeds per m². The three treatments were as follows: control (without application of folic acid), priming of seeds with 25 µM Solution of folic acid accompanying leaf spraying in steming stage. The third one was the same as second one except leaf spraying in heading stage. Folic acid spraying is carried out in the final hours of day by a pump sprayer. 100 mL of solution was sprayed per each square meter.

Finding: Results revealed that folic acid external application had following advantages compared to control: firstly, increase the yield of grain, Secondly, protein, essential amino acids, Iron content of grain is increased meaningfully and finally cause to increase the chlorophyll content in flag leaf.

Conclusions: In this research the effects of external application of folic acid have been studied on whole grain, but part of accumulated iron along with proteins during flour preparation process have been omitted. However considering positive impact of external application of folic acid on quality of grains and on proteins, amino acids, iron content of them; we can claim that this method was a short-term solution for both improving quality and quantity of food and reducing adverse effect of malnutrition.

Keywords: Iron, Phytic acid, Protein, Malnutrition, Zinc, Zn/PA ratio

*Corresponding author: esfandi1977@yahoo.com

Yield gap analysis of rice in relation to soil properties in Foumanat plain

N. Aghaeipour¹, *H. Pirdashti², M. Zavareh³, H. Asadi⁴ and M.A. Bahmanyar⁵

¹Ph.D. Student, Genetics and Agricultural Biotechnology Institute of Tabarestan, Sari Agricultural Sciences and Natural Resources University, Sari, Iran, ²Associate Prof., Physiology of Crops, Genetics and Agricultural Biotechnology Institute of Tabarestan, Sari Agricultural Sciences and Natural Resources University, Sari, Iran, ³Associate Prof., Physiology of Crops, University of Guilan, Rasht, Iran, ⁴Associate Prof., Dept., of Soil Science, University of Tehran, Iran, ⁵Professor, of Soil Science, Genetics and Agricultural Biotechnology Institute of Tabarestan, Sari Agricultural Sciences and Natural Resources University, Sari, Iran

Received: 15/03/2017; Accepted: 03/07/2017

Abstract

Background and objectives: Nowadays, due to population growth and changes in eating habits, there is a need to increase agricultural productions to meet consumer demand all around the world. Increasing the area under cultivation is one of the solutions for this problem, which cannot be appropriate in future because this needs to use of marginal lands with low yield and stability. Hence, increased yield is an effective strategy in this regard which can resolve the problem by decreasing yield gap. Considering the fact that there is an urgent need to increase rice production in the country as well as improving the productivity of soil resources, planning for the proper use of chemical fertilizers to achieve the maximum performance seems to be required. Accordingly, the aim of this study was to evaluate rice potential yield and yield gap in Foumanat plain using boundary line analysis, moreover, to find the relationship between soil properties and rice yield.

Materials and methods: In order to investigate the yield gap (YG) of rice (cv. 'Tarom Hashemi') related to soil properties, a field experiment was carried out in Foumanat plain during two cropping seasons: 2012-13 and 2013-14. We recorded the information of 53 fields as soil samples were taken and geographic coordinates were recorded. Some soil characteristics such as total nitrogen, potassium, phosphorus, organic matter and cation-exchange capacity were measured. At the end of growing season (harvesting time) grain yield was calculated in each field (1 m²). The boundary line method was used to calculate yield gap, optimum yield and optimum amount of soil properties for each field. Data analysis done by nonlinear regression of quadratic function, dent-like and segmented models based on PROC NLIN procedures.

Results: Segmented functions were well able to describe the trend of pH and electrical conductivity of soil. In addition, dent-like function used to describe the process of changes in available phosphorus, available potassium, organic matter and cation exchange capacity. Also, a quadratic function was used to describe the trend of total nitrogen. The average of optimum yield and actual yield were 7.67 and 4.81 ton ha⁻¹, respectively, in Foumanat plain with an YG of 2.86 ton ha⁻¹ (37.3 percent). The optimum concentration of total nitrogen, phosphorus, potassium, organic matter, cation exchange capacity, pH and electrical conductivity were 0.266 percent, 13.49-30.35 mg kg⁻¹, 102.2-173.4 mg kg⁻¹, 2.7-3.2 percent, 25.43-30.8 meq 100 g⁻¹, 6.36 and 1.92 dS m⁻¹, respectively.

Conclusion: Based on the results of this study, some of the main factors for rice YG in this area could be inappropriate management of chemical fertilizer usage (nitrogen, phosphorus and potassium) as well as ignored role of organic matter and cation exchange capacity in providing essential nutrients for rice. Boundary line analysis method could clearly calculate the soil properties potential in response to yield. In general, the results of this study can help to conclude a good strategy for achieving optimal production and gap reduction due to fertility situation of the area.

Keywords: Boundary line, Yield gap, Non-linear regression, Optimum yield, Soil resources

*Corresponding author: h.pirdashti@sanru.ac.ir

Study the effect of growth improver and chemical fertilizer application on growth and yield of *Solanum tuberosum*

*Z. Avarseji¹ and M. Tanha Khajeh²

¹Assistant Prof., Dept., of Plant Production, Gonbad Kavous University,

²B.Sc. Graduated of Plant Production

Received: 03/04/2017; Accepted: 18/09/2017

Abstract

Background and objectives: Potato is the fourth most important food crop in the world that plays an important role in people feeding. This crop is a source of carbohydrates, protein and essential amino acids for human being and it is one of the important tuber plants. Low yield of crops in many countries often related to the deficiency of soil nutrients. Chemical fertilizers have an important role in improving yield and quality of potatoes. However, excessive use of chemical fertilizers is associated with many health and environmental problems and pollution in water and soil. This study was conducted to evaluate the effect of organic fertilizers and growth promoters on potato (cultivar Milva) yield to reduce the consumption of chemical fertilizers.

Materials and methods: Experiment was conducted at the Agricultural Research Station of Gonbad Kavous University in a randomized complete block design with three replications in 2016. Treatments included the use of nitrogen, phosphorus and potassium to amount of common use of region (100 NPK), 100 NPK + Humic acid (HMC), 100 NPK + Equilibrate vegetative organic (EVEO), the use of nitrogen, phosphorus and potassium at a rate of half of the region convention (50 NPK) + HMC, 50 NPK + EVEO, application of EVEO and HMC alone. Traits of tuber number plant⁻¹, fresh and dry weight of tubers plant⁻¹, plant fresh weight ha⁻¹, biological yield, tuber yield, and harvest index were measured.

Results: the results indicated that application of HMC and EVEO with chemical fertilizers significantly increased tuber number plant⁻¹, tuber weight plant⁻¹, biological yield, tuber yield, and harvest index; but they had no significant effect on plant weight. The highest tuber number plant⁻¹, dry weight of tuber plant⁻¹, biological yield, tuber yield, and harvest index were obtained in 100 NPK + EVEO; although there was no significant difference with treatments of 100 NPK + HMC and 50 NPK + EVEO. The lowest amount of these traits were achieved in the treatment 100 NPK. However, it had no significant difference with treatments of application of EVEO and HMC alone.

Conclusion: Based on these results, the use of EVEO can reduce consumption of chemical fertilizers by 50%. Also, significant difference was not observed between chemical fertilizer treatment (100 NPK) and application of EVEO and HMC alone. Therefore, use of these organic materials can increase the absorption and efficiency of nutrition and consequently reduction of the consumption of chemical fertilizers will help to reduce environmental pollution.

Keywords: Chemical Fertilizer, Growth improver, Humic acid, Potato yield

*Corresponding author: avarseji@gonbad.ac.ir

Effect of potassium, zinc and boron foliar application on canopy temperature, physiological traits and yield of two bread wheat cultivars under optimum and late planting dates

H. Kamaei¹, *H.R. Eivand², M. Daneshvar³ and F. Nazarian⁴

¹Ph.D. Student of Agronomy, Lorestan University, ²Associate Prof., Dept., of Agronomy and Plant Breeding, Lorestan University, ³Assistant Prof., Dept., of Agronomy and Plant Breeding, Lorestan University, ⁴Professor Dept., of Agronomy and Plant Breeding, Lorestan University

Received: 09/04/2017; Accepted: 04/07/2017

Abstract

Background and objectives: wheat (*Triticum aestivum* L.) is one of the most important sources of plant food for human. High temperature resulting from delay in planting is one major environmental factor limiting growth and production of wheat, especially in tropical regions. One of the parameters that to evaluate the effect of heat stress in wheat genotypes used canopy temperature is. Most of the Iranian soils have high pH and calcareous nature. In this type of soils solubility of nutrients low, absorption of these elements by plant reduced and requirement of plants increases to these nutrients. This experiment was conducted to study the effect of potassium, zinc and boron foliar application on physiological traits and yield of two bread wheat cultivars under optimum and late planting dates.

Materials and Methods: An experiment was conducted as split-split plot based on randomized complete blocks design with 20 treatments and 3 replications in Ramhormoz city with latitude 31°16' N and longitude 49°37' E and a height of 160 meters above sea level in farming year 2015-16. The experimental factors were included planting date in two levels: optimum (November 21) and late (January 5) as the main factor, nutrients foliar application in five levels: (water as a control, potassium, zinc, boron and combination potassium + zinc + boron (each 3 lit.h⁻¹)) as the sub factor and two cultivars of bread wheat Pishtaz and Chamran 2 as the sub-sub factor. Solutions for foliar application were prepared by using potassium (21%), zinc-chelate (7.5%) and boron (5%). Traits measured were included leaf chlorophyll content, flag leaf relative water content, cell membrane stability index, maximum quantum efficiency of PSII (Fv/Fm) and leaf proline content.

Results: The results showed that with delay in planting all above traits except leaf proline content decreased significantly due to terminal heat stress. However, the delay in planting was increased canopy temperature. Potassium, zinc and boron foliar application separately and in combination reduced harmful effects caused by the delay in planting on the traits leaf chlorophyll content, flag leaf relative water content, cell membrane stability index, maximum quantum efficiency of PSII (Fv/Fm), leaf proline content and grain yield in both wheat cultivars. As well as, foliar application these nutrients reduced canopy temperature of wheat cultivars under both optimum and late planting dates. Despite the lack of significant differences in Pishtaz and Chamran 2 wheat cultivars in canopy temperature, cell membrane stability index, maximum quantum efficiency of PSII (Fv/Fm), leaf proline content and grain yield, these cultivars showed a more appropriate response to the zinc separate application and potassium + zinc + boron combined application in both optimum and late planting dates, respectively.

Conclusion: Chamran 2 wheat cultivar in conditions the lack of use nutrients foliar application of potassium, zinc and boron (control) showed more yield than Pishtaz cultivar under optimum and late planting dates, but with the use of nutrients foliar application of potassium, zinc and boron were different response bread wheat cultivars of Pishtaz and Chamran 2. Between canopy temperature with measured physiological traits and grain yield were observed significant relationship, so that loss of canopy temperature was improved physiological traits and subsequently increase yield bread wheat cultivars of Pishtaz and Chamran 2 under optimum and late planting dates, that this improvement was evident with the use of zinc foliar application for chamran 2 wheat cultivar and potassium+zinc+boron foliar application for Pishtaz wheat cultivar.

Keywords: Nutrients, Terminal heat stress

*Corresponding author: eivand.hr@lu.ac.ir